

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 365 364**  
**A2**

(12)

# EUROPEAN PATENT APPLICATION

(21) Application number: 89310857.1

(51) Int. Cl.<sup>5</sup>: **C07D 205/08**, **C07D 401/12**,  
**A61K 31/395**

(22) Date of filing: 20.10.89

(30) Priority: 20.10.88 JP 265183/88

(43) Date of publication of application:  
25.04.90 Bulletin 90/17

(84) Designated Contracting States:  
**AT BE CH DE FR GB IT LI LU NL SE**

(71) Applicant: **TAISHO PHARMACEUTICAL CO.**  
**LTD**  
**24-1 Takata 3-chome Toshima-ku**  
**Tokyo 171(JP)**


(72) Inventor: **Kawashima, Yutaka**  
**1731-1, Akodacho**  
**Tatebayashi-shi(JP)**  
Inventor: **Sato, Masakazu**

**15-1-6-205, Akamidai-2-chome**  
**Konosu-shi(JP)**  
Inventor: **Kawase, Masahiro**  
**Gurin Haitsu 102 1216-10, Kawarabuki**  
**Ageo-shi(JP)**  
Inventor: **Watanabe, Yoshiaki**  
**13-1-101, Ogawahigashicho-2-chome**  
**Kodaira-shi(JP)**  
Inventor: **Hatayama, Katsuo**  
**Danchi 35-3, 1200-215 Horisakicho**  
**Omiya-shi(JP)**


(74) Representative: **Lamb, John Baxter et al**  
**MARKS & CLERK 57/60 Lincoln's Inn Fields**  
**London WC2A 3LS(GB)**

(54) 2-Azetidinone derivatives.

(57) 2-Azetizinone derivatives, useful as blood platelet aggregation inhibiting agents, are those of the formula:




in which R<sup>1</sup> is a halogen atom a C<sub>1</sub>-C<sub>4</sub> alkyl group, a C<sub>1</sub>-C<sub>4</sub> alkoxy group or an alkoxycarbonyl group in which the alkoxy group is a C<sub>1</sub>-C<sub>4</sub> alkoxy group; R<sup>2</sup> is a group of the formula:




(in which R<sup>3</sup> and R<sup>4</sup> are the same or different and each is a hydrogen atom, a C<sub>1</sub>-C<sub>3</sub> alkenyl group, a

EP 0 365 364 A2


phenyl group or a benzyl group), a group of the formula:



(in which R<sup>5</sup> is a hydrogen atom, a phenyl group substituted by a halogen atom or a C<sub>1</sub>-C<sub>4</sub> alkoxy group, a phenyl group, a C<sub>1</sub>-C<sub>4</sub> alkyl group or a pyridyl group), a group of the formula:



(in which R<sup>6</sup> is a hydrogen atom, a C<sub>1</sub>-C<sub>4</sub> alkyl group or a benzyl group and R<sup>7</sup> is a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl group), a group of the formula:



(in which R<sup>8</sup> and R<sup>9</sup> are the same or different and each is a hydrogen atom or C<sub>1</sub>-C<sub>4</sub> alkyl group), a pyrrolidinyl group or a tetrahydroazepinyl group; and n is an integer of from 2 to 10; and salts thereof.


## 2-AZETIDINONE DERIVATIVES

The present invention relates to 2-azetidinone derivatives having blood platelet aggregation inhibiting activity.

Compounds with an azetidinone skeleton having blood platelet aggregation inhibiting activity are disclosed in Japanese Patent Kokai Nos. 62-87562, 62-87563, 63-225353 and 63-225354.

As a result of various researches, the present inventors have found novel 2-azetidinone derivatives having stronger blood platelet aggregation inhibiting activity, and have accomplished the present invention.

An object of the present invention is to provide a 2-azetidinone derivative represented by the formula:



wherein R<sup>1</sup> is a halogen atom, an alkyl group having 1 to 4 carbon atoms, an alkoxy group having 1 to 4 carbon atoms or an alkoxycarbonyl group in which the alkoxy group has 1 to 4 carbon atom, R<sup>2</sup> is a group represented by the formula




(wherein R<sup>3</sup> and R<sup>4</sup> are the same or different and are each a hydrogen atom, an alkyl group having 1 to 4 carbon atoms, an alkenyl group having 3 to 5 carbon atoms, a phenyl group or a benzyl group), a group represented by the formula



(wherein R<sup>5</sup> is a hydrogen atom, a phenyl group substituted by a halogen atom or an alkoxy group having 1 to 4 carbon atoms, a phenyl group, an alkyl group having 1 to 4 carbon atoms or a pyridyl group), a group represented by the formula



(wherein R<sup>6</sup> is a hydrogen atom, an alkyl group having 1 to 4 carbon atoms or a benzyl group, and R<sup>7</sup> is a hydrogen atom or an alkyl group having 1 to 4 carbon atoms), a group represented by the formula



(wherein  $R^8$  and  $R^9$  are the same or different and each a hydrogen atom or an alkyl group having 1 to 4 carbon atoms), a pyrrolidinyl group or a tetrahydroazepinyl group, and  $n$  is an integer of from 2 to 10), and a salt thereof.

## DETAILED DESCRIPTION OF THE INVENTION


In the present invention, the alkyl group having 1 to 4 carbon atoms refers to a straight or branched chain alkyl group such as a methyl group, an ethyl group, a propyl group, an isopropyl group and a butyl group. The alkoxy group having 1 to 4 carbon atoms refers to a straight or branched chain alkoxy group such as a methoxy group, an ethoxy group, a propoxy group, an isopropoxy group and a butoxy group. The alkenyl group having 3 to 5 carbon atoms may be an allyl group, a butenyl group, a prenyl group and the like. The halogen atoms may be a fluorine atom, a chlorine atom, a bromine atom or an iodine atom.

The salt means the pharmaceutically acceptable salt such as inorganic or organic acid salts (e.g., hydrochloride, sulfate, acetate, oxalate, maleate and the like).

Configuration of the iminoalkylidene substituent of the compound of the present invention is E-form, and configuration due to the asymmetric carbon atom at the 4-position is dl-form.

Among the preferred compounds of the present invention are (E)-1-(4-methoxyphenyl)-3-{2-[3-(4-morpholinyl)propoxyimino]propylidene}-4-phenyl-2-azetidinone and (E)-1-(4-methoxyphenyl)-3-[2-[3-{1-[4-(2-pyridyl)piperadinyll}propoxyimino]propylidene]-4-phenyl-2-azetidinone.

The compounds of the present invention can be prepared, for example, by the following method; a 2-azetidinone derivative represented by the formula



(II)

(wherein  $R^1$  is as defined above) is reacted with a hydroxylamine derivative represented by the formula  $H_2N-O-(CH_2)_n-R^2$  (III)

(wherein  $R^2$  and  $n$  are as defined above) in an inert solvent in the presence of a catalyst to give a compound of Formula I.


Examples of the inert solvent used herein are alcohols (e.g., methanol and ethanol), tetrahydrofuran, chloroform, methylene chloride, benzene, toluene, ethyl acetate, dioxane, xylene and the like. Examples of the catalyst are inorganic acids (e.g., hydrochloric acid gas and sulfuric acid) and the salts thereof, organic acids (e.g., p-toluenesulfonic acid, camphorsulfonic acid and acetic acid) and the salts thereof, amines (e.g., triethylamine and pyridine) and the salts thereof, and magnesium sulfate. The reaction temperature is from  $0^\circ C$  to the reflux temperature of the solvent, and preferably from room temperature to the reflux temperature of the solvent.

The compound of Formula II can be prepared, for example, by a process described in Japanese Patent Kokai 63-225354.

The compound of Formula III can be prepared, for example, by the following method; a compound represented by the formula

$Cl-(CH_2)_n-R^2$  (IV)

(wherein  $R^2$  and  $n$  are as defined above) is first reacted with N-hydroxylphthalimide in an inert solvent in the presence of a base to give a compound represented by the formula




(wherein  $R^2$  and  $n$  are as defined above).

15 Examples of the inert solvent are acetone, N,N-dimethylformamide and dimethyl sulfoxide, and examples of the base are sodium hydride, sodium amide, sodium hydroxide, potassium hydroxide, sodium carbonate, triethylamine and diazabicyclo[5,4,0]undec-7-ene. The reaction temperature is from 0°C to the reflux temperature of the solvent, and preferably from room temperature to the reflux temperature of the solvent.

20 The compound of Formula III are also obtained according to an ordinary manner, for example, by treating the compound of Formula V with hydrazine in an inert solvent as described above.

Alternatively, the compound of the present invention can be prepared as follows: 2-azetidinone represented by the formula



(wherein  $R^1$  is as defined above) is reacted with a compound of Formula IV in an inert solvent in the presence of a catalyst to give a compound of Formula I.

40 Examples of the inert solvent used herein are N,N-dimethylformamide, dimethyl sulfoxide, tetrahydrofuran. Examples of the catalyst are amines (e.g., triethylamine and diisopropylethylamine), salts (e.g., sodium carbonate and potassium carbonate) and sodium hydride. The reaction temperature is from 0°C to the reflux temperature of the solvent, and preferably from 0°C to the room temperature.

45 The compound of Formula VI can be prepared by reacting the compound of Formula II with hydroxylamine hydrochloride in an inert solvent in the presence of a catalyst.

Examples of the inert solvent used herein are alcohols (e.g., methanol, ethanol and isopropyl alcohol), tetrahydrofuran, chloroform, benzene and ethyl acetate. Examples of the catalyst are amines (e.g., imidazole and triethylamine), salts (e.g., potassium carbonate, sodium carbonate, sodium acetate and magnesium sulfate). The reaction temperature is from 0°C to the reflux temperature of the solvent, and preferably from 50 0°C to the room temperature.

The compound of the present invention have an excellent blood platelet aggregation inhibiting activity with very poor bleeding tendency as a side-effect, and therefore, they are useful as blood platelet aggregation inhibiting agent. For the purpose, these compounds can be administered orally or parenterally in a conventional dosage form such as tablets, powders, granules, capsules, solutions, suspensions, 55 injectional solutions and the like, each of which can be prepared by conventional pharmaceutical practices.

The dosage used as a blood platelet aggregation inhibiting agent to human depends on the age, weight, response of patient, administration route or time of administration, but usually it may be from 0.1 to 3000 mg per day.

The LD<sub>50</sub> of the compound of Formula I in mouse is more than 5000 mg/kg.

#### Experiment [Inhibitory effect on acute thrombocytopenia in mice]

Ten male ICR mice weighing 20 - 30 g were used for each group. Under pentobarbital anaesthesia, adenosine diphosphate (ADP) dissolved in physiological saline in a dose of 1 mg/kg was injected into the tail vein and 20  $\mu$ l of blood was collected from the femoral artery at 30 seconds after injection of ADP. Immediately after blood sampling, platelet counts were measured with an automatic blood cell counter (Sysmex CC-180A). Platelet count of normal group without injection of ADP was measured as well.

Test drugs [Compounds 2 and 3 (the compounds of the present invention) and Compound 1 (the comparative drug) in Table 1] were suspended in 5% gum arabic solution and administered by gavage in a dose of 300 mg/kg 2 hours prior to injection of ADP. As control group, 5% gum arabic solution without test drugs was administered.

Inhibition ratio of acute thrombocytopenia was calculated by the following formula.

$$\text{Inhibition ratio (\%)} = \frac{A - B}{C - B} \times 100$$

Note:

A: Platelet count of the group treated with both test drug and ADP

B: Platelet count of control group with both 5% gum arabic solution and ADP

C: Platelet count of normal group without injection of ADP

Results are shown in Table 1.

Table 1

Test drug	Inhibition Ratio (%)
Compound 1	12.00
Compound 2	54.66
Compound 3	57.61

Note:

Compound 1: (E)-1-(4-Methoxyphenyl)-3-(2-carboxymethoxyiminopropylidene)-4-phenyl-2-azetidinone

Compound 2: (E)-1-(4-Methoxyphenyl)-3-{2-[3-(4-morpholinyl)propoxyimino]propylidene}-4-phenyl-2-azetidinone

Compound 3: (E)-1-(4-Methoxyphenyl)-3- [ 2- [ 3-{1-[4-(2-pyridyl)piperadiny]-propoxyimino } -propylidene ] -4-phenyl-2-azetidinone

The present invention is illustrated by the following Reference Example and Examples in more detail.

#### Reference Example 1

To a suspension of 1.75 g of sodium hydride in 20 ml of N,N-dimethylformamide was added a mixture of 11.9 g of N-hydroxyphthalimide and 60 ml of N,N-dimethylformamide, and then the mixture was stirred for 30 minutes. To the reaction solution was added a mixture of 15 g of 3-chloropropylidibutylamine and 10

ml of N,N-dimethylformamide, and then the mixture was heated at reflux for 5 hours. After evaporation of the solvent, ethyl acetate was added to the residue, and the mixture was washed with water. Evaporation of the solvent gave 23.4 g of N-(3-dibutylaminopropoxy)phthalimide, which was then dissolved in 200 ml of methylene chloride. To the solution was added 20 ml of hydrazine monohydrate, and the mixture was stirred at room temperature for 3 hours. After removal of the insolubles by filtration, the filtrate was concentrated under reduced pressure and distilled under reduced pressure to give 10.3 g of O-(3-dibutylaminopropyl)hydroxylamine.  
b.p. 84 - 85 °C (0.5 mmHg)

Following a procedure similar to that of Reference Example 1, there were obtained the following compounds.  
O-[3-(4-Morpholinyl)propyl]-hydroxylamine.  
b.p. 80 - 82 °C (0.4 mmHg)  
O-[3-(Butylethylamino)propyl]-hydroxylamine.  
b.p. 68 - 70 °C (0.1 mmHg)  
O-[3-(4-Piperidyl)propyl]-hydroxylamine.  
b.p. 70 °C (0.6 mmHg)

#### Example 1

#### Preparation of (E)-1-phenyl-3-{2-[3-(1-piperidyl)propoxyimino]propylidene}-4-phenyl-2-azetidinone

A mixture of 1.11 g of (E)-1-phenyl-3-(2-oxo-propylidene)-4-phenyl-2-azetidinone, 0.63 g of O-[3-(1-piperidyl)propyl]-hydroxylamine, 46.5 mg of 10-camphorsulfonic acid and 40 ml of benzene was heated at reflux for 8 hours, and the solvent was evaporated under reduced pressure. The residue was chromatographed on alumina [eluent; hexane-acetone (7:3)] to give the fractions containing the end compound, and the solvent was evaporated. The residue was recrystallized from hexane to give 0.6 g of the title compound.  
m.p. 120 - 122 °C.

Following a procedure similar to that of Example 1, there were obtained the following compounds.  
(E)-1-(4-Fluorophenyl)-3-{2-[3-(1-piperidyl)propoxyimino]propylidene}-4-phenyl-2-azetidinone  
m.p. 118 - 120 °C  
(E)-1-(2-Methylphenyl)-3-{2-[3-(1-piperidyl)propoxyimino]propylidene}-4-phenyl-2-azetidinone oxalate  
<sup>1</sup>H - NMR (CDCl<sub>3</sub>) δ (ppm);  
1.44 - 1.75 (2H, m), 1.68 (3H, s), 1.80 - 2.22 (6H, m), 2.36 (3H, s), 2.70 - 3.40 (6H, m), 4.08 (2H, t), 5.75 (1H, d), 6.71 (1H, d), 6.98 - 7.20 (4H, m), 7.20 - 7.45 (5H, m), 8.00 (2H, brs)  
(E)-1-Phenyl-3-[2-(3-dibutylaminopropoxyimino)propylidene]-4-phenyl-2-azetidinone  
m.p. 59 - 61 °C  
(E)-1-(4-Fluorophenyl)-3-[2-(3-dibutylamino)propylidene]-4-phenyl-2-azetidinone  
m.p. 62 - 64 °C  
(E)-1-(2-Methylphenyl)-3-[2-(3-dibutylaminopropoxyimino)propylidene]-4-phenyl-2-azetidinone oxalate  
m.p. 69 - 71 °C  
(E)-1-Phenyl-3-[2-[3-(4-morpholinyl)propoxyimino]propylidene]-4-phenyl-2-azetidinone  
m.p. 87 - 89 °C  
(E)-1-(4-Fluorophenyl)-3-[2-[3-(4-morpholinyl)-propoxyimino]propylidene]-4-phenyl-2-azetidinone  
m.p. 121 - 123 °C  
(E)-1-(2-Methylphenyl)-3-[2-[3-(4-morpholinyl)-propoxyimino]propylidene]-4-phenyl-2-azetidinone  
m.p. 102 - 104 °C  
(E)-1-Phenyl-3-[2-(3-butylethylaminopropoxyimino)propylidene]-4-phenyl-2-azetidinone  
m.p. 76 - 79 °C  
(E)-1-(4-Fluorophenyl)-3-[2-(3-butylethylaminopropoxyimino)propylidene]-4-phenyl-2-azetidinone  
m.p. 76 - 79 °C  
(E)-1-(2-Methylphenyl)-3-[2-(3-butylethylaminopropoxyimino)propylidene]-4-phenyl-2-azetidinone oxalate  
<sup>1</sup>H - NMR (CDCl<sub>3</sub>) δ (ppm);  
0.93 (3H, t), 1.13 - 1.50 (5H, m), 1.50 - 1.80 (2H, m), 1.68 (3H, s), 1.92 - 2.20 (2H, m), 2.35 (3H, s), 2.90 - 3.30 (6H, m), 4.11 (2H, t), 5.77 (1H, d), 6.72 (1H, d), 7.00 - 7.23 (4H, m), 7.23 - 7.50 (5H, m), 8.95 (2H, brs)  
(E)-1-(4-Methoxyphenyl)-3-[2-[3-(4-morpholinyl)propoxyimino]propylidene]-4-phenyl-2-azetidinone  
(Compound 2)

- m.p. 106 - 108 °C  
 (E)-1-(4-Methoxyphenyl)-3-{2-[3-(1-piperidyl)propoxyimino]propylidene}-4-phenyl-2-azetidinone  
 m.p. 121 - 123 °C  
 (E)-1-(4-Methoxyphenyl)-3-{2-[3-(dibutylaminopropoxyimino)propylidene]-4-phenyl-2-azetidinone  
 5 m.p. 75 - 77 °C  
 (E)-1-(4-Methoxyphenyl)-3-{2-[3-(butylethylaminopropoxyimino)propylidene]-4-phenyl-2-azetidinone  
 m.p. 79 - 81 °C  
 (E)-1-(4-Methoxycarbonylphenyl)-3-{2-[3-(4-morpholinyl)propoxyimino]propylidene}-4-phenyl-2-azetidinone  
 m.p. 169 - 170.5 °C  
 10 (E)-1-(4-Methoxycarbonylphenyl)-3-{2-[3-(dibutylaminopropoxyimino)propylidene]-4-phenyl-2-azetidinone  
 m.p. 97 - 100 °C  
 (E)-1-(4-Methoxycarbonylphenyl)-3-{2-[3-(1-piperidyl)propoxyimino]propylidene}-4-phenyl-2-azetidinone  
 m.p. 187 - 170 °C  
 (E)-1-(4-Methoxycarbonylphenyl)-3-{2-[3-(butylethylaminopropoxyimino)propylidene]-4-phenyl-2-azetidinone  
 15 m.p. 95 - 97 °C

## Example 2

- 20 Preparation of (E)-1-(4-methoxyphenyl)-3-[2-[3-{1-[4-(2-pyridyl)piperadiny]-  
 }propoxyimino]propylidene]-4-phenyl-2-azetidinone (Compound 3)

(1) To a suspension of 9.2 g of (E)-1-(4-methoxyphenyl)-3-(2-oxopropylidene)-4-phenyl-2-azetidinone in  
 25 200 ml of isopropyl alcohol were added 3.1 g of hydroxylamine hydrochloride and 3.1 g of imidazole, and  
 then the mixture was stirred at room temperature overnight. The resulting precipitate was collected by  
 filtration, and recrystallized from isopropyl alcohol to give 8.8 g of (E)-1-(4-methoxyphenyl)-3-(2-hydrox-  
 ylaminopropylidene)-4-phenyl-2-azetidinone.  
 m.p. 204.5 - 206 °C.

- 30 (2) To a suspension of 0.4 g of sodium hydride in 20 ml of N,N-dimethylformamide was added a  
 mixture of 3.2 g of the compound obtained in the item (1) and 30 ml of N,N-dimethylformamide, and then  
 the mixture was stirred for 5 minutes. To the reaction mixture was added a mixture of 2.4 g of 1-(3-  
 chloropropyl)-4-(2-pyridyl)-piperadine and 20 ml of N,N-dimethylformamide, and the mixture was stirred at  
 35 room temperature overnight. To the reaction solution was added 200 ml of ethyl acetate, and the mixture  
 was washed 3 times with 200 ml of a saturated aqueous sodium chloride solution. After evaporation of the  
 solvent, the residue was recrystallized from a mixture of dichloromethane and hexane to give 2.6 g of the  
 title compound.  
 m.p. 153 - 154.5 °C

- Following a procedure similar to that of Example 2, there were obtained the following compounds.  
 40 (E)-1-(4-Methoxyphenyl)-3-[2-[3-[4-(3,5-dimethylmorpholinyl)]propoxyimino]propylidene]-4-phenyl-2-  
 azetidinone  
 m.p. 108 - 110 °C  
 (E)-1-(4-Methoxyphenyl)-3-[2-[3-[1-(4-benzylpiperidyl)]propoxyimino]propylidene]-4-phenyl-2-  
 azetidinone  
 45 m.p. 94.5 - 96.5 °C  
 (E)-1-(4-Methoxyphenyl)-3-[2-[3-(benzylethylaminopropoxyimino)propylidene]-4-phenyl-2-azetidinone  
 m.p. 81 - 83 °C  
 (E)-1-(4-Methoxyphenyl)-3-[2-[3-(diisobutylaminopropoxyimino)propylidene]-4-phenyl-2-azetidinone  
 m.p. 100 - 102 °C  
 50 (E)-1-(4-Methoxyphenyl)-3-[2-[3-(diallylaminopropoxyimino)propylidene]-4-phenyl-2-azetidinone  
 m.p. 87 - 89 °C  
 (E)-1-(4-Methoxyphenyl)-3-[2-[3-[1-(3,5-dimethylpiperidyl)]propoxyimino]propylidene]-4-phenyl-2-  
 azetidinone  
 m.p. 114.5 - 116.5 °C  
 55 (E)-1-(4-Methoxyphenyl)-3-[2-[3-(1-pyrrolidinyl)propoxyimino]propylidene]-4-phenyl-2-azetidinone  
 m.p. 110 - 112 °C  
 (E)-1-(4-Methoxyphenyl)-3-[2-[3-(1-tetrahydroazepinyl)propoxyimino]propylidene]-4-phenyl-2-azetidinone  
 m.p. 110 - 112 °C

- (E)-1-(4-Methoxyphenyl)-3-[2-(3-diethyl aminopropoxyimino)propylidene]-4-phenyl-2-azetidinone  
m.p. 90 - 92 °C
- (E)-1-(4-Methoxyphenyl)-3-[2-(3-benzylmethylaminopropoxyimino)propylidene]-4-phenyl-2-azetidinone  
m.p. 60 - 63 °C
- 5 (E)-1-(4-Methoxyphenyl)-3- [ 2-{3-[1-(4-methylpiperadiny)]propoxyimino}propylidene ] -4-phenyl-2-azetidinone  
m.p. 96 - 98 °C
- (E)-1-(4-Methoxyphenyl)-3-[2-(3-phenylmethylaminopropoxyimino)propylidene]-4-phenyl-2-azetidinone  
m.p. 122 - 124 °C
- 10 (E)-1-(4-Methoxyphenyl)-3-{2-[2-(4-morpholinyl)ethoxyimino]propylidene}-4-phenyl-2-azetidinone  
m.p. 96 - 98 °C
- (E)-1-(4-Methoxyphenyl)-3-{2-[2-(1-pyrrolidinyl)ethoxyimino]propylidene}-4-phenyl-2-azetidinone  
m.p. 103 - 105 °C
- (E)-1-(4-Methoxyphenyl)-3- [ 2- [ 3-{1-[4-(2-methoxyphenyl)piperadiny]- }propoxyimino ] propylidene ] -4-phenyl-2-azetidinone  
m.p. 126 - 128 °C
- 15 (E)-1-(4-Methoxyphenyl)-3- [ 2- [ 3-{1-[4-(4-fluorophenyl)piperadiny]]propoxyimino ] propylidene ] -4-phenyl-2-azetidinone  
m.p. 107 - 109 °C
- 20 (E)-1-(4-Methoxyphenyl)-3-[2-(2-benzylmethyl aminoethoxyimino)propylidene]-4-phenyl-2-azetidinone  
m.p. 93 - 95 °C
- (E)-1-(4-Methoxyphenyl)-3-{2-[6-(4-morpholinyl)hexyloxyimino]propylidene}-4-phenyl-2-azetidinone  
m.p. 86 - 88 °C
- (E)-1-(4-Methoxyphenyl)-3-{2-[6-(1-pyrrolidinyl)hexyloxyimino]propylidene}-4-phenyl-2-azetidinone  
25 m.p. 99 - 101 °C
- (E)-1-(4-Methoxyphenyl)-3-[2-(6-benzylmethylaminohexyloxyimino)propylidene]-4-phenyl-2-azetidinone  
m.p. 73 - 75 °C
- (E)-1-(4-Methoxyphenyl)-3-[2-(6-butylethylaminohexyloxyimino)propylidene]-4-phenyl-2-azetidinone  
m.p. 73 - 75 °C
- 30


### Claims

1. 2-Azetidinone derivatives of the formula:

35

40


45



50

in which R<sup>1</sup> is a halogen atom, a C<sub>1</sub>-C<sub>4</sub> alkyl group, a C<sub>1</sub>-C<sub>4</sub> alkoxy group or an alkoxycarbonyl group in which the alkoxy group is a C<sub>1</sub>-C<sub>4</sub> alkoxy group; R<sup>2</sup> is a group of the formula:


55




(In which R<sup>3</sup> and R<sup>4</sup> are the same or different and each is a hydrogen atom, a C<sub>1</sub>-C<sub>5</sub> alkenyl group, a phenyl group or a benzyl group), a group of the formula:



(in which R<sup>5</sup> is a hydrogen atom, a phenyl group substituted by a halogen atom or a C<sub>1</sub>-C<sub>4</sub> alkoxy group, a phenyl group, a C<sub>1</sub>-C<sub>4</sub> alkyl group or a pyridyl group), a group of the formula:



(in which R<sup>6</sup> is a hydrogen atom, a C<sub>1</sub>-C<sub>4</sub> alkyl group or a benzyl group, and R<sup>7</sup> is a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl group), a group of the formula:



(in which R<sup>8</sup> and R<sup>9</sup> are the same or different and each is a hydrogen atom or C<sub>1</sub>-C<sub>4</sub> alkyl group), a pyrrolidinyl group or a tetrahydroazepinyl group; and n is an integer of from 2 to 10; and salts thereof.

2. (E)-1-(4-methoxyphenyl)-3{2-[3-(4-morpholinyl)propoxyimino]propylidene}-4-phenyl-2-azetidinone.

3. (E)-1-(4-methoxyphenyl)-3-2-[3-{1-[4-(2-pyridyl)piperidinyl]propoxyimino]propylidene]-4-phenyl-2-azetidinone.

4. A pharmaceutical composition comprising an azetidinone derivative as claimed in claim 1 in association with a pharmaceutical carrier or diluent.



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



Publication number:

**0 365 364 A3**

12

# EUROPEAN PATENT APPLICATION

21 Application number: 89310857.1

51 Int. Cl.<sup>5</sup>: **C07D 205/10, C07D 401/12, A61K 31/395**

22 Date of filing: 20.10.89

30 Priority: 20.10.88 JP 265183/88

43 Date of publication of application:  
25.04.90 Bulletin 90/17

84 Designated Contracting States:  
**AT BE CH DE FR GB IT LI LU NL SE**

86 Date of deferred publication of the search report:  
11.12.91 Bulletin 91/50

71 Applicant: **TAISHO PHARMACEUTICAL CO. LTD**  
24-1 Takata 3-chome Toshima-ku  
Tokyo 171(JP)

72 Inventor: Kawashima, Yutaka  
1731-1, Akodacho

**Tatebayashi-shi(JP)**

Inventor: **Sato, Masakazu**  
15-1-6-205, Akamidai-2-chome  
**Konosu-shi(JP)**

Inventor: **Kawase, Masahiro**  
Gurin Hatsu 102 1216-10, Kawarabuki  
**Ageo-shi(JP)**


Inventor: **Watanabe, Yoshiaki**  
13-1-101, Ogawahigashicho-2-chome  
**Kodaira-shi(JP)**

Inventor: **Hatayama, Katsuo**  
Danchi 35-3, 1200-215 Horisakicho  
**Omiya-shi(JP)**


74 Representative: **Lamb, John Baxter et al**  
**MARKS & CLERK** 57-60 Lincoln's Inn Fields  
London WC2A 3LS(GB)

54 **2-Azetidinone derivatives.**


57 2-Azetidinone derivatives, useful as blood platelet aggregation inhibiting agents, are those of the formula:




in which R<sup>1</sup> is a halogen atom a C<sub>1</sub>-C<sub>4</sub> alkyl group, a C<sub>1</sub>-C<sub>4</sub> alkoxy group or an alkoxy carbonyl group in which the alkoxy group is a C<sub>1</sub>-C<sub>4</sub> alkoxy group; R<sup>2</sup> is a group of the formula:




(in which R<sup>3</sup> and R<sup>4</sup> are the same or different and each is a hydrogen atom, a C<sub>1</sub>-C<sub>5</sub> alkenyl group, a phenyl group or a benzyl group), a group of the formula:



(in which R<sup>5</sup> is a hydrogen atom, a phenyl group substituted by a halogen atom or a C<sub>1</sub>-C<sub>4</sub> alkoxy group, a phenyl group, a C<sub>1</sub>-C<sub>4</sub> alkyl group or a pyridyl group), a group of the formula:



(in which R<sup>6</sup> is a hydrogen atom, a C<sub>1</sub>-C<sub>4</sub> alkyl group or a benzyl group and R<sup>7</sup> is a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl group), a group of the formula:



(in which R<sup>8</sup> and R<sup>9</sup> are the same or different and each is a hydrogen atom or C<sub>1</sub>-C<sub>4</sub> alkyl group), a pyrrolidinyl group or a tetrahydroazepinyl group; and n is an integer of from 2 to 10; and salts thereof.



European  
Patent Office

## EUROPEAN SEARCH REPORT

Application Number

EP 89 31 0857

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
Y	EP-A-0 264 232 (TAISHO PHARMACEUTICAL CO.) * Claims * & JP-A-63 225 354 (Cat. D) - - -	1,4	C 07 D 205/10 C 07 D 401/12 A 61 K 31/395
A	US-A-4 647 457 (S. ADAM-MOLINA) * Claims * - - -	1,4	
P,Y	CHEMICAL ABSTRACTS, vol. 113, no. 13, 24th September 1990, page 669, abstract no. 115062r, Columbus, Ohio, US; & JP-A-01 246 256 (TAISHO PHARMACEUTICAL CO., LTD) 02-10-1989 * Abstract * - - - - -	1,4	
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			C 07 D 205/00 C 07 D 401/00 C 07 D 499/00 A 61 K 31/00
The present search report has been drawn up for all claims			
Place of search The Hague		Date of completion of search 01 October 91	Examiner CHOULY J.
<div>CATEGORY OF CITED DOCUMENTS</div> <div>X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention</div> <div>E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons ----- &amp;: member of the same patent family, corresponding document</div>			

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☒ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**